

RESISTANCE PATTERN OF EXTENDED DRUG RESISTANT SALMONELLA TYPHI IN CHILDREN

ANSWAR¹, SALAHUDDIN A², AHMAD A^{1*}

1. Pediatrics Department HMC/ KGMC, Peshawar, Pakistan
 2. Pediatrics Department, Peshawar General Hospital Peshawar Pakistan
- *Corresponding author email address: ambreenahmad1971@yahoo.com

(Received, 15th April 2023, Revised 7th October 2023, Published 9th October 2023)

Abstract To determine the frequency of antibiotic resistance of extended drug-resistant *Salmonella typhi* in children. Cross-sectional observational study was conducted at Pediatric B Unit, Hayatabad Medical Complex, Peshawar, from 1st April 2021 to 30th June 2022. Approval from the institutional ethical committee was obtained. Using nonprobability consecutive sampling technique, this study was conducted on 104 pediatric patients admitted to Children B Unit with blood culture-proven enteric fever. Data was collected from blood cultures and sensitivity reports from the microbiology section of our hospital laboratory. This study was conducted on 32 female and 72 male pediatric in-patients with blood culture-proven enteric fever caused by *Salmonella typhi*. Resistance to Azithromycin was 43.26%, Meropenem 17.30%, Polymixin 13.46%, Colistin 31.73%, Tigecycline, 01.92%, Imipenem 04.80% *Salmonella typhi* demonstrated high levels of resistance to commonly used antibiotics.

Keywords: Antibiotic resistance, *Salmonella typhi*, children, enteric fever

Introduction

Enteric fever is considered a significant health issue in developing countries where risk factors for contracting this infectious disease, like poverty, overpopulation, lack of clean drinking water and sanitation, and improper disposal of human waste, are prevalent. Moreover, indiscriminate and irrational use of antimicrobial agents is rampant. This has resulted in widespread resistance to various antimicrobial treatments for enteric fever. Reports of the emergence of extensive antibiotic resistance among isolates of *Salmonella typhi* from all over Pakistan are appearing in literature (M. Khan et al., 2022). Pediatric age group is also affected by drug-resistant *Salmonella typhi* (S. Ahmad et al., 2020; F. Z. Khan, Baig, Zameer, & Sharafat, 2018).

This increasing spread of resistance among pathogenic strains of *Salmonella typhi* has resulted in the loss of affordable antibiotic options for the treatment of enteric fever, compelling clinicians to use costly antibiotics like azithromycin and carbapenems (Memon, Ahmed, & Iqbal, 2020; Umair & Siddiqui, 2020). Indiscriminate use of these relatively newer agents may result in resistance. This may lead to complete treatment failure if the trend of increasing resistance is not stopped (Hussain et al., 2019). This fact necessitates surveillance of antibiotic resistance of *Salmonella typhi* by using blood culture to detect resistance and choose the effective antibiotics (Anjum et al., 2021; Hameed et al., 2019). Antibiotic stewardship and selective use

of antibiotics, according to sensitivity report, is essential to deal with this challenge (Chaudhry, 2022; Nusrat et al., 2022). This study was undertaken to determine the resistance pattern of *Salmonella typhi* strains causing enteric fever in our patients.

Materials and methods

Material & Methods: Approval from the institutional ethical committee was obtained and using non probability consecutive sampling technique, this study was conducted on 104 pediatric patients admitted to Children B Unit, Hayatabad Medical Complex Peshawar. Inclusion criteria were pediatric patients with clinical features consistent with enteric fever and isolation of *Salmonella typhi* from blood culture. Exclusion criteria were clinical features of acute febrile illnesses like malaria, urinary tract infection, meningitis, etc. Data was collected from blood culture and sensitivity reports from the microbiology section of our hospital laboratory using a proforma prepared for data collection. Results were entered in Microsoft Excel 2010 spreadsheet and analyzed for calculating percentages. Results were presented in tables.

Results: This study was conducted on 32 female and 72 male pediatric in-patients with blood culture-proven enteric fever caused by *Salmonella typhi*. Resistance to azithromycin was 43.26%, meropenem 17.30%, polymixin 13.46%, colistin 31.73%, tigecycline, 01.92%, imipenem 04.80%

Table 1: Antibiotic Resistance Pattern of Salmonella Typhi

S No	Antibiotic	Resistant	Percentage
1	azithromycin	45	43.26%
2	meropenem	18	17.30%
3	polymixin	14	13.46%
4	colistin	33	31.73%
5	tigecycline	02	01.92%
6	imipenem	05	04.80%

Discussion

Our study shows 45 isolates out of 104 (43.26%) resistant to azithromycin. Literature reporting rising resistance of Salmonella typhi against conventional antibiotics and newer agents like azithromycin, carbapenems, and tigecycline is a genuinely alarming phenomenon (Butt et al., 2022). Salmonella isolates carrying mutations that confers azithromycin resistance have been reported (Iqbal et al., 2020). A study by Ahmad reported 55.81% resistance to azithromycin (A. Ahmad et al., 2021). 93 out of 323 (28.7%) isolates were resistant to azithromycin studied by Shaikh and colleagues (Shaikh, Shaikh, & Tahir, 2019). A study from Islamabad in 322 patients found 204 (63.4%) isolates azithromycin resistant (Umair & Siddiqui, 2020). A study from Rawalpindi showed 62 out of 108 (57.4%) isolates resistant to azithromycin. The findings of these studies are supporting one another and may be pointing to the presence and progress of resistance to azithromycin (Menz et al., 2021).

This study detected resistance to meropenem in 18 (17.30%). 47.7% isolates were resistant to meropenem in a study by Shah SA et al. (Shah et al., 2020) Out of 108 isolates 44 (40.7%) were reported resistant to meropenem by Shah SA and colleagues (Browne et al., 2020). These findings indicate a grave situation if such a high level of resistance spread because it will make Salmonella typhi infection incurable.

Our study detected 05/104 (04.80%) Salmonella typhi isolates showing resistance to imipenem. In comparison, Amber reported 1.8% resistance (Amber, Fatima, Qurat-ul-ain Khalid, Khan, & Zulfqar, 2021). Other studies had reported no resistance in their patients (Malik & Ahmed, 2016; Yasin et al., 2022). Our study showed resistance of 31.73% to colistin, 13.46 % to polymixin B, and 1.92% to tigecycline. Local data showing recent trends in resistance to polymixins and tigecycline almost non-existent.

Conclusions

Salmonella typhi demonstrated high levels of resistance to commonly used antibiotics. This indicates a strong need for judicious antibiotic use based on culture and sensitivity reports of the isolated strain of Salmonella typhi.

References

- Ahmad, A., Mustaan, S., Wahab, F., Ayyaz, M., Sadiq, A., & Hussain, I. (2021). Pattern of Antibiotic Resistance in Salmonella Typhi Isolates with Special Consideration to Extended Drug Resistant Typhoid. *Journal of Saidu Medical College, Swat*, **11**(2), 103-107.
- Ahmad, S., Zahid, S. B., Salahuddin, A., Khan, A., Khan, M. H., & Ali, H. M. (2020). Trends in antibiotic susceptibility of enteric fever isolates among children attending a tertiary care hospital of Peshawar, KP. *Journal of Rehman Medical Institute*, **6**(3), 20-23.
- Amber, W., Fatima, I., Qurat-ul-ain Khalid, H. H., Khan, I. M., & Zulfqar, R. (2021). Antibiotic Resistance Pattern of Salmonella Species in Children in a Tertiary Care Hospital. *Journal of Rawalpindi Medical College*, **25**(3).
- Anjum, M., Soomro, S., Kulsoom, S., Bibil, S., Asim, S., & Riaz, M. (2021). Clinical Spectrum, Laboratory Profile and Antibiotic Susceptibility Pattern of Children with Enteric Fever at a Tertiary Care Hospital of Karachi. *Journal of Islamic International Medical College (JIIMC)*, **16**(1), 4-9.
- Browne, A. J., Kashef Hamadani, B. H., Kumaran, E. A., Rao, P., Longbottom, J., Harriss, E., . . . Baker, S. (2020). Drug-resistant enteric fever worldwide, 1990 to 2018: a systematic review and meta-analysis. *BMC Medicine*, **18**, 1-22.
- Butt, M. H., Saleem, A., Javed, S. O., Ullah, I., Rehman, M. U., Islam, N., . . . Misbah, S. (2022). Rising XDR-typhoid fever cases in Pakistan: are we heading back to the pre-antibiotic era? *Frontiers in public health*, **9**, 794868.
- Chaudhry, A. R. (2022). *Clinical Spectrum, Bacteriological Profile and Antibiogram of Culture Proven Enteric Fever in a Private Sector Hospital in Lahore*. Paper presented at the Medicinsk Forum.
- Hameed, S., Izhar, M., Basheer, A., Lal, C., Rishi, S., & Basit, A. (2019). An Update on Isolation of Extensively Drug Resistant (XDR) Salmonella enterica from Blood Cultures in a Tertiary Care Centre. *Proceedings SZPGMI Vol*, **33**(4), 20-24.
- Hussain, A., Satti, L., Hanif, F., Zehra, N. M., Nadeem, S., Bangash, T. M., & Peter, A. (2019). Typhoidal Salmonella strains in Pakistan: an impending threat of extensively drug-resistant Salmonella Typhi. *European Journal of Clinical Microbiology & Infectious Diseases*, **38**, 2145-2149.
- Iqbal, J., Dehraj, I. F., Carey, M. E., Dyson, Z. A., Garrett, D., Seidman, J. C., . . . Qamar, F. N. (2020). A race against time: reduced azithromycin susceptibility in Salmonella

- enterica serovar Typhi in Pakistan. *mSphere*, **5**(4), 10.1128/msphere.00215-00220.
- Khan, F. Z., Baig, S., Zameer, S., & Sharafat, S. (2018). Emerging Trends of Antibiotic Resistance pattern of Salmonella Typhi. *International Journal of Pathology*, 52-57.
- Khan, M., Tu-Zahra, F., Gul, A., Saleem, M., Riaz, M., & Gul, W. (2022). Extensively Drug-Resistant Salmonella enterica serovar Typhi in Pediatric population during CoVID-19 Pandemic. *International Journal of Infectious Diseases*, **116**, S4.
- Malik, N., & Ahmed, M. (2016). In vitro effect of new antibiotics against clinical isolates of Salmonella Typhi. *Journal of the College of Physicians and Surgeons--Pakistan*, **26**(4), 288-292.
- Memon, A., Ahmed, M., & Iqbal, K. (2020). Frequency and antimicrobial resistance pattern of extensive-drug resistance Salmonella typhi isolates. *Journal of Microbiology and Infectious Diseases*, **10**(02), 68-74.
- Menz, B. D., Charani, E., Gordon, D. L., Leather, A. J., Moonesinghe, S. R., & Phillips, C. J. (2021). Surgical antibiotic prophylaxis in an era of antibiotic resistance: common resistant bacteria and wider considerations for practice. *Infect Drug Resist*, 5235-5252.
- Nusrat, N., Islam, M. R., Paul, N., Rahman, N., Krishnapillai, A., Haq, M. A., & Haque, M. (2022). Clinical and Laboratory Features of Enteric Fever in Children and Antibiotic Sensitivity Pattern in a Tertiary Care Hospital of a Low-and Middle-Income Country. *Cureus*, **14**(10).
- Shah, S. A. A., Nadeem, M., Syed, S. A., Abidi, S. T. F., Khan, N., & Bano, N. (2020). Antimicrobial sensitivity pattern of Salmonella Typhi: emergence of resistant strains. *Cureus*, **12**(11).
- Shaikh, A. A., Shaikh, A., & Tahir, A. (2019). Antimicrobial resistance trends of typhoidal Salmonellae in southern Pakistan. *Rawal Medical Journal*, **44**(1), 7-7.
- Umair, M., & Siddiqui, S. A. (2020). Antibiotic susceptibility patterns of Salmonella typhi and Salmonella paratyphi in a tertiary care hospital in Islamabad. *Cureus*, **12**(9).
- Yasin, N., Rahman, H., Sarwar, Y., Qasim, M., Nisa, I., Ikram, A., . . . Khan, N. (2022). Salmonella Typhi from Northwest Pakistan: Molecular strain typing and drug resistance signature. *Microbial Drug Resistance*, **28**(1), 120-126.

Declarations

Data Availability statement

All data generated or analyzed during the study are included in the manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Funding

Not applicable

Conflict of Interest

Regarding conflicts of interest, the authors state that their research was carried out independently without any affiliations or financial ties that could raise concerns about biases.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2023