

## BIOLUMINESCENCE SYSTEMS

IDREES N<sup>1\*</sup>, SIDDIQUE A<sup>1</sup>, FATEH A<sup>1</sup>, AFZAL N<sup>2</sup>, ALI I<sup>3</sup>, JAVIED MA<sup>1\*</sup>, ALI Q<sup>1</sup>, MALIK A<sup>1</sup>

<sup>1</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Pakistan

<sup>2</sup>Department of Biotechnology, Virtual University of Pakistan

<sup>3</sup>Department of Microbiology and Molecular Genetics, University of Okara, Pakistan

First six authors equally contribute towards write up and final editing of manuscript

\*Corresponding author email: [falakawais7@gmail.com](mailto:falakawais7@gmail.com), [nimijutt1998@gmail.com](mailto:nimijutt1998@gmail.com)

(Received, 07<sup>th</sup> August 2020, Revised 25<sup>th</sup> November 2020, Published 30<sup>th</sup> November 2020)

**Abstract:** *Bioluminescence System has been mostly studied in reporter technology. There are forty different bioluminescent systems occurred in nature while there are only seven of these light emitting systems and their biochemical reactions have been studied however the pathway of only two biochemical systems has been understood yet. Here, we have provided an overview for these bioluminescent systems as a tool for researchers working on bioluminescence for better future applications of bioluminescent.*

**Keywords:** Bioluminescence, Luciferase, Luciferin, synthetic biology and imaging

### Introduction

The bioluminescence gives living creatures an unmistakable preferred position in certain natural settings. The capacity to transmit lights under darkness conditions has been reported around in 10,000 species out of 800 genera, despite the fact that it likely could be an irony (Chen *et al.*, 2012; Haddock *et al.*, 2010). The specific advantage from light discharges in under different conditions has long way from being to clear for various species, under any condition, by and large bioluminescence has been thought to fill the need of seeing or visualizing correspondence to frighten away from the predators, draw in prey and or in romance conduct (Baker *et al.*, 2019; Ellis and Oakley, 2016; Janicke *et al.*, 2018; Wainwright and Longo, 2017). Developments have unearthed and have fixed various answers for bioluminescence biochemical process exhibiting that capacity to gleam has always been available only to the living organisms for different purposes of even form microorganisms to higher animals like insects (ladybird battle). Different luciferins, the little particles involved in inclined to light outflows upon the oxidation biochemical reactions, which have been brought from increase in inconsequential biochemical reaction pathways (Kotlobay *et al.*, 2018; Roda and Guardigli, 2012). The oxidation of the atoms are usually catalyzed through non-homologous types of compounds, *i.e.*, luciferases, to make a smooth light-producing responses which are usually diverse in shading, the rate of catalysis, cell localization and reliance on NADH, ATP and different secondary

metabolites are involved (Kaskova *et al.*, 2016; Li *et al.*, 2018; Vacher *et al.*, 2018). Although it has been still generally understood on sub-atomic and atomic levels, the scarcely could such be assorted in variety of the responses in an effectively quantifiable yield turning into a basic piece of present day correspondent innovations. Luminescent responses, where structure for both of the luciferases and luciferin have been reported found, which are currently used for food testing (Carling and Huang, 2013; Huang *et al.*, 2015; Shama and Malik, 2013), ecological observing and identifications (Girotti *et al.*, 2008; Lima *et al.*, 2011), diagnostics studies (Burakova *et al.*, 2015; Frank and Krasitskaya, 2014), tranquilize screenings (Hasson *et al.*, 2015; Sun *et al.*, 2016) and different sorts of biomedical and clinical explorations. There have been discovered various types of luciferins Luciferases (Kotlobay *et al.*, 2018), the biology of bioluminescence mechanisms (Hall *et al.*, 2012; Widder, 2010) which are considered as just as an extensive outline for all of the known bioluminescent frameworks (Dixon *et al.*, 2016; Xing *et al.*, 2016) and which are easily accessible. Various researchers have given very purposefully and concise outline of light-transmitting responses where both of the luciferase and luciferin are known, featuring for their principle highlights for reasonable applications. There have been assembled a large number of bioluminescent frameworks while keeping their structures of luciferins as marker identifications and these mixes the vital determinants for shading as well as the characteristics of light-discharging responses in any of the event form forty

bioluminescent frameworks which have been discovered and still exist in nature in different forms (Cannaert *et al.*, 2016; Yao *et al.*, 2018).

#### **Coelenterazine-subordinate structure**

The finest arranged for the structures of bioluminescent has been found in marine animals. In marine water, there is a regular and comparable coelenterazine and luciferin which fills in with substrates for different unreservedly progressed luciferases in phylogenetically blocked off the social affairs among living creatures of marine ecosystem (Haddock *et al.*, 2001; Teranishi and Shimomura, 1997). Coelenterazine is a change tripeptide conveyed from a single phenylalanine and two tyrosine molecules, regardless, the characteristics shown during its biosynthesis, similarly as the particular biosynthetic way, and characteristics shown in dark condition (Shimomura and Teranishi, 2000; Yao *et al.*, 2018). Most of marine living creatures don't consolidate coelenterazine themselves; rather, they gain it from the food likely clarification behind an earth shattering centered headway for bioluminescence under marine conditions. The coelenterazine-subordinate structures under natural conditions emanates blue light, of about 450–500 nm, which don't need any cofactors for enzymes except need for oxygen. From time to time, the shade of bioluminescence changes by a fluorescent protein which is associated with luciferases (Inouye and Shimomura, 1997; Mosrin *et al.*, 2009). Various qualities, like sub-nuclear weight, pH-affectability, thermo-stability, and catalysis space for luciferases usually move radically among the coelenterazine-subordinate bioluminescent systems. There are two noteworthy luciferases which utilizes coelenterazine or its analogs. The renilla luciferases: a medium-sized of about 36 kDa cytoplasmic protein found in corals which makes a fast sharp signal. The early divergence (Contag and Bachmann, 2002; Lorenz *et al.*, 1991), similarly, the availability for the planned transformations along with an extended magnificence and red-moved spectrum (Berg *et al.*, 2009; Islam *et al.*, 2012; Wilson and Hastings, 1998), made its structure notable for applications in biomedical especially, in drug screening and bioimaging (Newman *et al.*, 2011; Regot *et al.*, 2014). The gaussia luciferases: have a size about 20 kDa free cytoplasmic protein which released by a small shellfish for social affair Copepoda, which with high rate of catalysis and remarkable thermo stability. The formation of these proteins depends upon the formation and improvement of disulfide bonds which makes it differ for various heterologous protein structures. Indication of gaussia luciferases was found scaled straightly with the cells number being analyzed during study (James and Gambhir, 2012;

Skog *et al.*, 2008; Tannous *et al.*, 2005) which makes this system more significant for analyzing the drug responses and tumor development. The nanoluc luciferases: a structured variety of luciferases found and isolated from the shrimp *Oplophorus gracilirostris*. This small (19 kDa) protein utilizes a selective permeable coelenterazine, it needs disulfide bonds. Mixes up with fluorescent producing proteins which achieve a splendid structure bioluminescent form with red-moved spectra empowering single-cell and whole-body bioluminescent imaging in vivo (Chu *et al.*, 2016; Rangaraju *et al.*, 2014; Saito *et al.*, 2012).

#### **Cypridina luciferin-based bioluminescent system**

The cypridina luciferin is a tripeptide molecule which emits blue light during luminescence, which is produced due to a metabolite found in ostracod *Cypridina* which is a bioluminescent marine fish, also called as *Porichthys notatus*. The isoleucine, tryptophan and arginine are usually causes of biochemical reactions for bioluminescence (Oba *et al.*, 2002; Vacher *et al.*, 2018). The *Cypridina* bioluminescent systems have been found to be used commonly in bioimaging, immunoassays (Kaskova *et al.*, 2016; Strassburg *et al.*, 2012; Wu *et al.*, 2009) and for the examinations of circadian rhythms (Kaskova *et al.*, 2016; Yamada *et al.*, 2013).

#### **D-Luciferin Dependent System**

Luciferin system deals with the rules of light transmitting reactions and determinants of colors. Luciferase protein has been extricated from firefly which creates light within the sight of substrate luciferin. Oxygen and ATP act a source of energy that converts d-luciferin into oxyluciferin so light is delivered by utilizing the synthetic energy (Shimomura, 2006; Tannous *et al.*, 2005). The additional linkages of beetles, click creepy crawlies, fireflies and rail streets as they are for all intents and purposes significant for bioluminescent response as they are stable and non-toxic compound. They have the property of discharging light in a detached framework with various colors and ranges (blue having wide range spectrum peaking at 480nm), green (508nm) with limited range and there are numerous colors too, like red and orange which represent independent origin of a similar bioluminescent system (Dixon *et al.*, 2016; Widder, 2010). Enzymes, for example, aminoacyl tRNA synthetase and acetyl-coA ligases are canalized by a cofactor of Mg<sup>+</sup> and utilize ATP as an energy packet with a protein, so D-luciferin gets oxidized by luciferase enzyme. The light omitted by them relies on the convergence of ATP, so that the system produce by the light fills in as ATP indicator and malignancy of metabolism indicator through monetizing microorganisms contaminating in water

and to check the degree of ATP in the blood (Cannaert *et al.*, 2016; Huang *et al.*, 2015; Islam *et al.*, 2012).

Firefly luciferase catalyst changes over to luciferin, because of its property of creating glow it is utilized in pharmaceuticals, bioanalytical applications. It is highly recommended because of its high quantum yield of bioluminescence, accessibility of thermo stability, mutant varieties and improved qualities of easily creation in microbes and viruses (Frank and Krasitskaya, 2014; Oba *et al.*, 2002). The accessibility of protein ligand trade to examine metabolites implicated in cell signaling and its correspondence, among different applications. It is additionally utilized as a research particle having high quantum yield of bioluminescence. Click beetle luciferase; D-luciferin subordinate luciferase get from pyro phosphorus plagiotalamus is a multi-colored heteroprotein complementation system empowered by double shading capability of two discrete sets of collaborating proteins (Xing *et al.*, 2016; Yao *et al.*, 2018). It is second mainstream gathering of D-Luciferin dependent system. The emission of light has been found by four sorts of luciferase with emanation of greatest range green light 540nm-593nm orange – red. They are commercially designed with variation for instance Chroma-luciferase offered by promega, they offer shading assortment, resistance to wide scope of PH, makes it appealing for numerous applications (Inouye and Shimomura, 1997; Rangaraju *et al.*, 2014; Vacher *et al.*, 2018).

#### **Meganyctiphanes Norvegica SARs**

The huge part of sparkles shown on the outside of sea is because of dinoflagellates creating shine or because of unicellular Algae, Dinoflagellates. Bioluminescence is limited in exceptional organelles like scintillators which happen with flashes of light accelerated by electrical and mechanical stimulations (Kartvedt, 2010; Schmidt, 2010; Spicer and Saborowski, 2010). It goes about as a protection instrument making their attacks obvious which is looking for consideration from predators; from high tropic level. The luciferin in these events are tetrapyrrole containing four to five piece of one nitrogen and four carbon compounds and its oxidation is catalyzed by dinoflagellates, luciferase achieves blue green light engaged at around 470nm (Saito *et al.*, 2012; Shama and Malik, 2013). Dinoflagellates and euphemistic krill uses two comparable tetrapyrrole based luciferins and structure another huge gathering of bioluminescent species. They are also utilized in research as an apparatus of inaccessibility of engineered luciferin (Cleary *et al.*, 2012; Yang *et al.*, 2002).

#### **Bacterial bioluminescent frame**

All bioluminescent microscopic organisms uses a similar special component for light discharge, where photons are created in responses requiring oxygen, flavin mononucleotide, myristic aldehyde and nicotinamide adenine dinucleotide. Over the span of responses, myristic aldehyde is oxidized and is subsequently known as luciferin, despite the fact that the genuine light source in bacterial bioluminescence is flavin mononucleotide. Bacterial luciferases are comprises of two polypeptide chains which structures a complex (75 kDa) and are encoded in the lux operon along with catalysts catalyzing luciferin biosynthesis. As a rule, bioluminescence is blue (~490 nm), in any case, both characteristic (Contag *et al.*, 1997; Daubner *et al.*, 1987) and designed (Ke and Tu, 2011; Xu *et al.*, 2016) red-moved forms of the bacterial frameworks. The full pathway for luciferin biosynthesis has been known since late 80's making lux operon the main hereditarily encodable bioluminescent accessible over the most recent thirty years (Bhuckory *et al.*, 2019; Hwang *et al.*, 2019). The flavin mononucleotide, myristic aldehyde and nicotinamide adenine dinucleotide are the main cause of gleaming living beings, including microorganisms (Belas *et al.*, 1982; Meighen, 1991), mammalian cell lines (Patterson *et al.*, 2005; Singh *et al.*, 2008), yeasts (Gupta *et al.*, 2003; Nivens *et al.*, 2004), plants (Daniell *et al.*, 2016; Krichevsky *et al.*, 2010) and bacteria and fungi. Nonetheless, the bioluminescent multicellular life forms have been made, maybe because of harmfulness or shortcoming of the mechanisms in eukaryotes (Burakova *et al.*, 2015). Among the primary utilizations of the mechanisms are the investigations of antimicrobial medications and bacterial contaminations (Belas *et al.*, 1982; Wu *et al.*, 2009). The most brilliant form to produce bioluminescence created to date is Lux (Brodl *et al.*, 2018; Gregor *et al.*, 2018).

#### **Fungal bioluminescent system**

A biochemical pathway creating bioluminescence has been portrayed totally, giving the principle innately encodable pathway from eukaryotes (Kotlobay *et al.*, 2018; Shimomura, 2006; Slot and Gluck-Thaler, 2019; Zhou *et al.*, 2020). The biochemical pathway utilizes an essential  $\alpha$ -pyrone 3-hydroxyhispidin which is oxidized by an insoluble luciferase in a biochemical reaction which just requires oxygen and results in the transmission of green light (~520 nm). A wild-type *Neonothopanus nambi* luciferase, nnLuz, is valuable in an arrangement of heterologous structures, with the introduction like that of the firefly luciferase (Roda and Guardigli, 2012; Zhou *et al.*, 2020). It has been demonstrated that the affirmation of these three characteristics from the infectious bioluminescent system is satisfactory to fabricate eukaryotes.

## Conclusion

In the presence of bioluminescent systems which is reasonable for various assessments and applications, diverse light-produces responses which involves various specialties in present-day correspondence advancements, In bioimaging, utilization of bioluminescence and fluorescence-based methodologies cover, a high unique range, low foundation, or profound tissue imaging. Precautionary measures and investigations of bacterial science are regularly finding the bacterial bioluminescent system while sedate screenings frequently utilizes D-luciferin-subordinate system. While choosing a luciferin-luciferase pair for a specific application, a few models must be considered including thermo-stability, ideal pH, protein size, inner and outer cell area, accumulation properties, outflow frequency, force, the space for the response reliance on ATP, and different cofactors. An ongoing revelation of a eukaryote-accommodating hereditarily encodable pathway in growth may animate the improvement of new bioluminescence-based innovation that would not need an expansion of the substrate. The capability of a bioluminescence-based device in manufactured science has been just barely investigated. The field of bioluminescence is shockingly under study. Simultaneously, with new bits of knowledge in photophysics, heredity qualities, and nature of bioluminescence being made each year, building new light-producing and light-conveying living system is getting more open than any time in recent advances.

## Conflict of interest

The authors declared absence of any conflict of interest.

## References

- Baker, L. J., Freed, L. L., Easson, C. G., Lopez, J. V., Fenolio, D., Sutton, T. T., Nyholm, S. V., and Hendry, T. A. (2019). Diverse deep-sea anglerfishes share a genetically reduced luminous symbiont that is acquired from the environment. *Elife* **8**, e47606.
- Belas, R., Mileham, A., Cohn, D., Hilman, M., Simon, M., and Silverman, M. (1982). Bacterial bioluminescence: isolation and expression of the luciferase genes from *Vibrio harveyi*. *Science* **218**, 791-793.
- Berg, J., Hung, Y. P., and Yellen, G. (2009). A genetically encoded fluorescent reporter of ATP: ADP ratio. *Nature methods* **6**, 161-166.
- Bhuckory, S., Kays, J. C., and Dennis, A. M. (2019). In Vivo Biosensing Using Resonance Energy Transfer. *Biosensors* **9**, 76.
- Brodl, E., Winkler, A., and Macheroux, P. (2018). Molecular mechanisms of bacterial bioluminescence. *Computational and structural biotechnology journal* **16**, 551-564.
- Burakova, L. P., Kudryavtsev, A. N., Stepanyuk, G. A., Baykov, I. K., Morozova, V. V., Tikunova, N. V., Dubova, M. A., Lyapustin, V. N., Yakimenko, V. V., and Frank, L. A. (2015). Bioluminescent detection probe for tick-borne encephalitis virus immunoassay. *Analytical and bioanalytical chemistry* **407**, 5417-5423.
- Cannaert, A., Storme, J., Franz, F., Auwärter, V., and Stove, C. P. (2016). Detection and activity profiling of synthetic cannabinoids and their metabolites with a newly developed bioassay. *Analytical chemistry* **88**, 11476-11485.
- Carling, P. C., and Huang, S. S. (2013). Improving healthcare environmental cleaning and disinfection current and evolving issues. *Infection Control & Hospital Epidemiology* **34**, 507-513.
- Chen, Y., Spiering, A., Karthikeyan, S., Peters, G. W., Meijer, E., and Sijbesma, R. P. (2012). Mechanically induced chemiluminescence from polymers incorporating a 1, 2-dioxetane unit in the main chain. *Nature chemistry* **4**, 559-562.
- Chu, J., Oh, Y., Sens, A., Ataie, N., Dana, H., Macklin, J. J., Laviv, T., Welf, E. S., Dean, K. M., and Zhang, F. (2016). A bright cyan-excitable orange fluorescent protein facilitates dual-emission microscopy and enhances bioluminescence imaging in vivo. *Nature biotechnology* **34**, 760-767.
- Cleary, A. C., Durbin, E. G., and Rynearson, T. A. (2012). Krill feeding on sediment in the Gulf of Maine (North Atlantic). *Marine Ecology Progress Series* **455**, 157-172.
- Contag, C. H., and Bachmann, M. H. (2002). Advances in in vivo bioluminescence imaging of gene expression. *Annual review of biomedical engineering* **4**, 235-260.
- Contag, C. H., Contag, P. R., and Benaron, D. A. (1997). Non-invasive localization of a light-emitting conjugate in a mammal. Google Patents.
- Daniell, H., Lin, C.-S., Yu, M., and Chang, W.-J. (2016). Chloroplast genomes: diversity, evolution, and applications in genetic engineering. *Genome biology* **17**, 134.
- Daubner, S. C., Astorga, A. M., Leisman, G. B., and Baldwin, T. O. (1987). Yellow light emission of *Vibrio fischeri* strain Y-1: purification and characterization of the energy-accepting yellow fluorescent protein. *Proceedings of the National Academy of Sciences* **84**, 8912-8916.
- Dixon, A. S., Schwinn, M. K., Hall, M. P., Zimmerman, K., Otto, P., Lubben, T. H., Butler, B. L., Binkowski, B. F., Machleidt, T.,

- and Kirkland, T. A. (2016). NanoLuc complementation reporter optimized for accurate measurement of protein interactions in cells. *ACS chemical biology* **11**, 400-408.
- Ellis, E. A., and Oakley, T. H. (2016). High rates of species accumulation in animals with bioluminescent courtship displays. *Current Biology* **26**, 1916-1921.
- Frank, L. A., and Krasitskaya, V. V. (2014). Application of enzyme bioluminescence for medical diagnostics. In "Bioluminescence: Fundamentals and Applications in Biotechnology-Volume 1", pp. 175-197. Springer.
- Girotti, S., Ferri, E. N., Fumo, M. G., and Maiolini, E. (2008). Monitoring of environmental pollutants by bioluminescent bacteria. *Analytica chimica acta* **608**, 2-29.
- Gregor, C., Gwosch, K. C., Sahl, S. J., and Hell, S. W. (2018). Strongly enhanced bacterial bioluminescence with the lux operon for single-cell imaging. *Proceedings of the National Academy of Sciences* **115**, 962-967.
- Gupta, R. K., Patterson, S. S., Ripp, S., Simpson, M. L., and Saylor, G. S. (2003). Expression of the *Photobacterium luminescens* lux genes (luxA, B, C, D, and E) in *Saccharomyces cerevisiae*. *FEMS yeast research* **4**, 305-313.
- Haddock, S. H., Moline, M. A., and Case, J. F. (2010). Bioluminescence in the sea. *Annual review of marine science* **2**, 443-493.
- Haddock, S. H., Rivers, T. J., and Robison, B. H. (2001). Can coelenterates make coelenterazine? Dietary requirement for luciferin in cnidarian bioluminescence. *Proceedings of the National Academy of Sciences* **98**, 11148-11151.
- Hall, M. P., Unch, J., Binkowski, B. F., Valley, M. P., Butler, B. L., Wood, M. G., Otto, P., Zimmerman, K., Vidugiris, G., and Machleidt, T. (2012). Engineered luciferase reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. *ACS chemical biology* **7**, 1848-1857.
- Hasson, S. A., Fogel, A. I., Wang, C., MacArthur, R., Guha, R., Heman-Ackah, S., Martin, S., Youle, R. J., and Inglese, J. (2015). Chemogenomic profiling of endogenous PARK2 expression using a genome-edited coincidence reporter. *ACS chemical biology* **10**, 1188-1197.
- Huang, Y.-S., Chen, Y.-C., Chen, M.-L., Cheng, A., Hung, I.-C., Wang, J.-T., Sheng, W.-H., and Chang, S.-C. (2015). Comparing visual inspection, aerobic colony counts, and adenosine triphosphate bioluminescence assay for evaluating surface cleanliness at a medical center. *American journal of infection control* **43**, 882-886.
- Hwang, E., Song, J., and Zhang, J. (2019). Integration of nanomaterials and bioluminescence resonance energy transfer techniques for sensing biomolecules. *Biosensors* **9**, 42.
- Inouye, S., and Shimomura, O. (1997). The Use of Renilla Luciferase, Oplophorus Luciferase, and Apoaequorin as Bioluminescent Reporter Protein in the Presence of Coelenterazine Analogues as Substrate. *Biochemical and biophysical research communications* **233**, 349-353.
- Islam, M. N., Das, S. R., Emin, M. T., Wei, M., Sun, L., Westphalen, K., Rowlands, D. J., Quadri, S. K., Bhattacharya, S., and Bhattacharya, J. (2012). Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nature medicine* **18**, 759-765.
- James, M. L., and Gambhir, S. S. (2012). A molecular imaging primer: modalities, imaging agents, and applications. *Physiological reviews* **92**, 897-965.
- Janicke, T., Ritchie, M. G., Morrow, E. H., and Marie-Orleach, L. (2018). Sexual selection predicts species richness across the animal kingdom. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20180173.
- Kaartvedt, S. (2010). Diel vertical migration behaviour of the northern krill (*Meganyctiphanes norvegica* Sars). In "Advances in Marine Biology", Vol. 57, pp. 255-275. Elsevier.
- Kaskova, Z. M., Tsarkova, A. S., and Yampolsky, I. V. (2016). 1001 lights: luciferins, luciferases, their mechanisms of action and applications in chemical analysis, biology and medicine. *Chemical Society Reviews* **45**, 6048-6077.
- Ke, D., and Tu, S. C. (2011). Activities, kinetics and emission spectra of bacterial luciferase-fluorescent protein fusion enzymes. *Photochemistry and photobiology* **87**, 1346-1353.
- Kotlobay, A. A., Sarkisyan, K. S., Mokrushina, Y. A., Marcet-Houben, M., Serebrovskaya, E. O., Markina, N. M., Somermeyer, L. G., Gorokhovatsky, A. Y., Vvedensky, A., and Purtov, K. V. (2018). Genetically encodable bioluminescent system from fungi. *Proceedings of the National Academy of Sciences* **115**, 12728-12732.
- Krichevsky, A., Meyers, B., Vainstein, A., Maliga, P., and Citovsky, V. (2010). Autoluminescent plants. *PLoS one* **5**, e15461.

- Li, J.-B., Chen, L., Wang, Q., Liu, H.-W., Hu, X.-X., Yuan, L., and Zhang, X.-B. (2018). A bioluminescent probe for imaging endogenous peroxynitrite in living cells and mice. *Analytical chemistry* **90**, 4167-4173.
- Lima, T. M., Procópio, L. C., Brandão, F. D., Leão, B. A., Tótola, M. R., and Borges, A. C. (2011). Evaluation of bacterial surfactant toxicity towards petroleum degrading microorganisms. *Bioresource technology* **102**, 2957-2964.
- Lorenz, W. W., McCann, R. O., Longiaru, M., and Cormier, M. J. (1991). Isolation and expression of a cDNA encoding Renilla reniformis luciferase. *Proceedings of the National Academy of Sciences* **88**, 4438-4442.
- Meighen, E. A. (1991). Molecular biology of bacterial bioluminescence. *Microbiology and Molecular Biology Reviews* **55**, 123-142.
- Mosrin, M., Bresser, T., and Knochel, P. (2009). Regio- and chemoselective multiple functionalization of chloropyrazine derivatives. Application to the synthesis of coelenterazine. *Organic Letters* **11**, 3406-3409.
- Newman, R. H., Fosbrink, M. D., and Zhang, J. (2011). Genetically encodable fluorescent biosensors for tracking signaling dynamics in living cells. *Chemical reviews* **111**, 3614-3666.
- Nivens, D. E., McKnight, T., Moser, S., Osbourn, S., Simpson, M., and Sayler, G. (2004). Bioluminescent bioreporter integrated circuits: potentially small, rugged and inexpensive whole-cell biosensors for remote environmental monitoring. *Journal of Applied Microbiology* **96**, 33-46.
- Oba, Y., Kato, S.-i., Ojika, M., and Inouye, S. (2002). Biosynthesis of luciferin in the sea firefly, *Cypridina hilgendorffii*: L-tryptophan is a component in *Cypridina* luciferin. *Tetrahedron letters* **43**, 2389-2392.
- Patterson, S. S., Dionisi, H. M., Gupta, R. K., and Sayler, G. S. (2005). Codon optimization of bacterial luciferase (lux) for expression in mammalian cells. *Journal of industrial microbiology and biotechnology* **32**, 115-123.
- Rangaraju, V., Calloway, N., and Ryan, T. A. (2014). Activity-driven local ATP synthesis is required for synaptic function. *Cell* **156**, 825-835.
- Regot, S., Hughey, J. J., Bajar, B. T., Carrasco, S., and Covert, M. W. (2014). High-sensitivity measurements of multiple kinase activities in live single cells. *Cell* **157**, 1724-1734.
- Roda, A., and Guardigli, M. (2012). Analytical chemiluminescence and bioluminescence: latest achievements and new horizons. *Analytical and bioanalytical chemistry* **402**, 69-76.
- Saito, K., Chang, Y., Horikawa, K., Hatsugai, N., Higuchi, Y., Hashida, M., Yoshida, Y., Matsuda, T., Arai, Y., and Nagai, T. (2012). Luminescent proteins for high-speed single-cell and whole-body imaging. *Nature communications* **3**, 1-9.
- Schmidt, K. (2010). Food and feeding in Northern krill (*Meganyctiphanes norvegica* Sars). In "Advances in marine biology", Vol. 57, pp. 127-171. Elsevier.
- Shama, G., and Malik, D. J. (2013). The uses and abuses of rapid bioluminescence-based ATP assays. *International journal of hygiene and environmental health* **216**, 115-125.
- Shimomura, O. (2006). "Bioluminescence: chemical principles and methods," World Scientific.
- Shimomura, O., and Teranishi, K. (2000). Light-emitters involved in the luminescence of coelenterazine. *Luminescence: the journal of biological and chemical luminescence* **15**, 51-58.
- Singh, H., Manuri, P. R., Olivares, S., Dara, N., Dawson, M. J., Huls, H., Hackett, P. B., Kohn, D. B., Shpall, E. J., and Champlin, R. E. (2008). Redirecting specificity of T-cell populations for CD19 using the Sleeping Beauty system. *Cancer research* **68**, 2961-2971.
- Skog, J., Würdinger, T., Van Rijn, S., Meijer, D. H., Gainche, L., Curry, W. T., Carter, B. S., Krichevsky, A. M., and Breakefield, X. O. (2008). Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nature cell biology* **10**, 1470-1476.
- Slot, J. C., and Gluck-Thaler, E. (2019). Metabolic gene clusters, fungal diversity, and the generation of accessory functions. *Current Opinion in Genetics & Development* **58**, 17-24.
- Spicer, J. I., and Saborowski, R. (2010). Physiology and metabolism of Northern krill (*Meganyctiphanes norvegica* Sars). In "Advances in marine biology", Vol. 57, pp. 91-126. Elsevier.
- Strassburg, K., Huijbrechts, A. M., Kortekaas, K. A., Lindeman, J. H., Pedersen, T. L., Dane, A., Berger, R., Brenkman, A., Hankemeier, T., and van Duynhoven, J. (2012). Quantitative profiling of oxylipins through comprehensive LC-MS/MS analysis: application in cardiac surgery. *Analytical and bioanalytical chemistry* **404**, 1413-1426.
- Sun, N., Youle, R. J., and Finkel, T. (2016). The mitochondrial basis of aging. *Molecular cell* **61**, 654-666.
- Tannous, B. A., Kim, D.-E., Fernandez, J. L., Weissleder, R., and Breakefield, X. O. (2005).

- Codon-optimized Gaussia luciferase cDNA for mammalian gene expression in culture and in vivo. *Molecular Therapy* **11**, 435-443.
- Teranishi, K., and Shimomura, O. (1997). Coelenterazine analogs as chemiluminescent probe for superoxide anion. *Analytical biochemistry* **249**, 37-43.
- Vacher, M., Fdez. Galván, I., Ding, B.-W., Schramm, S., Berraud-Pache, R., Naumov, P., Ferré, N., Liu, Y.-J., Navizet, I., and Roca-Sanjuán, D. (2018). Chemi- and bioluminescence of cyclic peroxides. *Chemical reviews* **118**, 6927-6974.
- Wainwright, P. C., and Longo, S. J. (2017). Functional innovations and the conquest of the oceans by acanthomorph fishes. *Current Biology* **27**, R550-R557.
- Widder, E. A. (2010). Bioluminescence in the ocean: origins of biological, chemical, and ecological diversity. *Science* **328**, 704-708.
- Wilson, T., and Hastings, J. W. (1998). Bioluminescence. *Annual review of cell and developmental biology* **14**, 197-230.
- Wu, C., Irie, S., Yamamoto, S., and Ohmiya, Y. (2009). A bioluminescent enzyme immunoassay for prostaglandin E2 using Cypridina luciferase. *Luminescence* **24**, 131-133.
- Xing, S., Wallmeroth, N., Berendzen, K. W., and Grefen, C. (2016). Techniques for the analysis of protein-protein interactions in vivo. *Plant physiology* **171**, 727-758.
- Xu, T., Close, D., Handagama, W., Marr, E., Sayler, G., and Ripp, S. (2016). The expanding toolbox of in vivo bioluminescent imaging. *Frontiers in oncology* **6**, 150.
- Yamada, Y., Nishide, S.-y., Nakajima, Y., Watanabe, T., Ohmiya, Y., Honma, K.-i., and Honma, S. (2013). Monitoring circadian time in rat plasma using a secreted Cypridina luciferase reporter. *Analytical biochemistry* **439**, 80-87.
- Yang, N.-C., Ho, W.-M., Chen, Y.-H., and Hu, M.-L. (2002). A convenient one-step extraction of cellular ATP using boiling water for the luciferin-luciferase assay of ATP. *Analytical biochemistry* **306**, 323-327.
- Yao, Z., Zhang, B. S., and Prescher, J. A. (2018). Advances in bioluminescence imaging: new probes from old recipes. *Current opinion in chemical biology* **45**, 148-156.
- Zhou, X., Mehta, S., and Zhang, J. (2020). Genetically encodable fluorescent and bioluminescent biosensors light up signaling networks. *Trends in Biochemical Sciences*.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).